




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 417/14, A61K 31/40 C07D 413/14	A1	(11) International Publication Number: WO 93/18032 (43) International Publication Date: 16 September 1993 (16.09.93)
(21) International Application Number: PCT/US93/01667 (22) International Filing Date: 3 March 1993 (03.03.93) (30) Priority data: 846,640 5 March 1992 (05.03.92) US (60) Parent Application or Grant (63) Related by Continuation US 846,640 (CIP) Filed on 5 March 1992 (05.03.92) (71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).		(72) Inventors; and (75) Inventors/Applicants (for US only) : MACOR, John, E. [US/US]; 42 Ann Avenue, Mystic, CT 06355 (US). NOWAKOWSKI, Jolanta, T. [US/US]; 37 Wood Cart Lane. Haddam, CT 06438 (US). (74) Agents: RICHARDSON, Peter, C. et al.; Pfizer Inc., Patent Department, 235 East 42nd Street, New York, NY 10017 (US). (81) Designated States: AU, BR, CA, CZ, DE (Utility model), JP, KR, NO, NZ, PL, RU, SK, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report.

(54) Title: INDOLE DERIVATIVES AS 5-HT₁ AGONISTS

(57) Abstract

Compounds of formula (I) where A represents a direct bond, C₁-C₄ alkyl, or C₁-C₄ alkenyl; n is 0, 1, or 2; R₁ is hydrogen, C₁-C₆ alkyl, aryl, C₁-C₃ alkylaryl, C₁-C₃ alkylheteroaryl, or -(CH₂)_mR₆; W, X, Y, and Z are each independently oxygen, sulfur, nitrogen or carbon, provided that at least one of W, X, Y or Z is nitrogen; R₂, R₃, R₄, and R₅ are each independently hydrogen, C₁-C₆ alkyl, aryl, C₁-C₃ alkylaryl, C₁-C₃ alkylheteroaryl, halogen, cyano, trifluoromethyl, nitro, -OR₇, -NR₇R₈, -(CH₂)_sOR₇, -SR₇, -SO₂NR₇R₈, -NR₇SO₂R₈, -NR₇CO₂R₈, -CONR₇R₈, or -CO₂R₇; one of R₂ and R₃, R₃ and R₄, or R₄ and R₅ may be taken together to form a five- to seven-membered alkyl ring, a six-membered aryl ring, a five- to seven-membered heteroalkyl ring having 1 heteroatom of N, O, or S, or a five- to six-membered heteroaryl ring having 1 or 2 heteroatoms of N, O, or S; R₆ is cyano, trifluoromethyl, or -OR₉; R₇, R₈, and R₉ are each independently hydrogen, C₁ to C₆ alkyl, -(CH₂)_mR₁₀, C₁ to C₃ alkylaryl, or aryl; R₇ and R₈ may be taken together to form a C₄-C₇ alkyl ring; R₁₀ is cyano, trifluoromethyl, or C₁-C₄ alkoxy; R₁₁ is hydrogen, -OR₁₂, or -NHCOR₁₂; R₁₂ is C₁ to C₆ alkyl, aryl, or C₁ to C₃ alkyl-aryl; m is 1, 2, or 3; s is 0, 1, 2, or 3; and the above aryl groups and the aryl moieties of the above alkylaryl groups are independently selected from phenyl and substituted phenyl, wherein said substituted phenyl may be substituted with one to three groups selected from C₁ to C₄ alkyl, halogen, hydroxy, cyano, carboxamido, nitro, and C₁ to C₄ alkoxy, and the pharmaceutically acceptable salts thereof. These compounds are useful psychotherapeutics and are potent serotonin (5-HT₁) agonists and may be used in the treatment of depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain and chronic paroxysmal hemicrania and headache associated with vascular disorders, and other disorders arising from deficient serotonergic neurotransmission. The compounds can also be used as centrally acting antihypertensives and vasodilators.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

5

INDOLE DERIVATIVES AS 5-HT₁ AGONISTSBackground of the Invention

The present invention relates to indole derivatives, to processes and intermediates for their preparation, to pharmaceutical compositions containing them and to their medicinal use. The active compounds of the present invention are useful in
10 treating migraine and other disorders.

United States Patents 4,839,377 and 4,855,314 and European Patent Application Publication Number 313397 refer to 5-substituted 3-aminoalkyl indoles. The compounds are said to be useful for the treatment of migraine.

British Patent Application 040279 refers to 3-aminoalkyl-1H-indole-5-thioamides
15 and carboxamides. The compounds are said to be useful in treating hypertension, Raymond's disease and migraine.


European Patent Application Publication Number 303506 refers to 3-poly:hydro-pyridyl-5-substituted-1H-indoles. The compounds are said to have 5-HT₁ receptor agonist and vasoconstrictor activity and to be useful in treating migraine.

20 European Patent Application Publication Number 35477 refers to N-piperidiny:indolyl:ethyl-alkane sulfonamide derivatives. The compounds are said to have 5-HT₁ receptor agonist and vasoconstrictor activity and to be useful in treating cephalic pain.

European Patent Application Publication Numbers 438230, 494774, and 497512
25 refer to indole-substituted five-membered heteroaromatic compounds. The compounds are said to have 5-HT₁-like receptor agonist activity and to be useful in the treatment of migraine and other disorders for which a selective agonist of these receptors is indicated.

Summary of the Invention

30 The present invention relates to compounds of the formula



35

-2-

where A represents a direct bond, C₁-C₄ alkyl, or C₁-C₄ alkenyl; n is 0, 1, or 2; R₁ is hydrogen, C₁-C₆ alkyl, aryl, C₁-C₃ alkylaryl, C₁-C₃ alkylheteroaryl, or -(CH₂)_mR₆; W, X, Y, and Z are each independently oxygen, sulfur, nitrogen or carbon, provided that at least one of W, X, Y, or Z is nitrogen; R₂, R₃, R₄, and R₅ are each independently
5 hydrogen, C₁-C₆ alkyl, aryl, C₁-C₃ alkylaryl, C₁-C₃ alkylheteroaryl, halogen, cyano, trifluoromethyl, nitro, -OR₇, -NR₇R₈, -(CH₂)_sOR₇, -SR₇, -SO₂NR₇R₈, -NR₇SO₂R₈, -NR₇CO₂R₈, -CONR₇R₈, or -CO₂R₇; one of R₂ and R₃, R₃ and R₄, or R₄ and R₅ may be taken together to form a five- to seven-membered alkyl ring, a six-membered aryl ring, a five- to seven-
10 membered heteroalkyl ring having 1 heteroatom of N, O, or S, or a five- to six-membered heteroaryl ring having 1 or 2 heteroatoms of N, O or S; R₆ is cyano, trifluoromethyl, or -OR₉; R₇, R₈, and R₉ are each independently hydrogen, C₁ to C₆ alkyl, -(CH₂)_mR₁₀, C₁ to C₃ alkylaryl, or aryl; R₇ and R₈ may be taken together to form a C₄-C₇ alkyl ring; R₁₀ is cyano, trifluoromethyl, or C₁-C₄ alkoxy; R₁₁ is hydrogen, -OR₁₂, or -NHCOR₁₂; R₁₂ is C₁ to C₆ alkyl, aryl, or C₁ to C₃ alkyl-aryl; m is 1, 2, or 3; s is 0, 1, 2,
15 or 3; and the above aryl groups and the aryl moieties of the above alkylaryl groups are independently selected from phenyl and substituted phenyl, wherein said substituted phenyl may be substituted with one to three groups selected from C₁ to C₄ alkyl, halogen (e.g. fluorine, chlorine, bromine or iodine), hydroxy, cyano, carboxamido, nitro, and C₁ to C₄ alkoxy, and the pharmaceutically acceptable salts thereof. These
20 compounds are useful in treating migraine and other disorders.

The compounds of the invention include all optical isomers of formula I (e.g., R and S stereogenicity at any chiral site) and their racemic, diastereomeric, or epimeric mixtures. When R₁₁ is hydrogen, the epimers with the R absolute configuration at the chiral carbon site designated by an asterisk in formula I are preferred. When R₁₁ is
25 -OR₁₂ or -NHCOR₁₂ and n is 0 or 1, the epimers with the S absolute configuration at the chiral carbon site designated by an asterisk in formula I are preferred. When R₁₁ is -OR₁₂ or -NHCOR₁₂ and n is 2, the epimers with the R absolute configuration at the chiral carbon site designated by an asterisk in formula I are preferred. When R₁₁ is -OR₁₂ or -NHCOR₁₂ and n is 0, the cis epimers [(2S, 3S) absolute configuration in the
30 azetidine ring] are particularly preferred. When R₁₁ is -OR₁₂ or -NHCOR₁₂ and n is 1, the cis epimers [(2S, 4R) absolute configuration in the pyrrolidine ring] are particularly preferred. When R₁₁ is -OR₁₂ or -NHCOR₁₂ and n is 2, the cis epimers [(2R, 5R) absolute configuration in the piperidine ring] are particularly preferred.

-3-

Unless otherwise indicated, the alkyl groups referred to herein, as well as the alkyl moieties of other groups referred to herein (e.g. alkoxy), may be linear or branched, and they may also be cyclic (e.g. cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl) or be linear or branched and contain cyclic moieties.

5 Preferred compounds of the invention are compounds of the formula I wherein A is either a direct bond or $-\text{CH}_2-$; n is 1; R_1 is hydrogen, C_1 - C_4 alkyl or $-\text{CH}_2\text{CH}_2\text{OCH}_3$; Z is nitrogen; Y is carbon; W and X are each independently oxygen, sulfur, nitrogen or carbon; and R_2 , R_3 , and R_4 are as defined above. Of the foregoing preferred compounds, when R_{11} is hydrogen, the epimers with the R absolute configuration at the
10 chiral carbon site designated by an asterisk in formula I are more preferred. Of the foregoing preferred compounds, when R_{11} is $-\text{OR}_{12}$ or $-\text{NHCOR}_{12}$, the epimers with the S absolute configuration at the chiral carbon site designated by an asterisk in formula I are more preferred. Of the foregoing compounds, when R_{11} is $-\text{OR}_{12}$ or $-\text{NHCOR}_{12}$, the cis epimers [(2S, 4R) absolute configuration in the pyrrolidine ring] are
15 particularly preferred.

The following compounds are particularly preferred:

(R)-5-(4-benzyl-1,3-thiazol-2-yl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;
(R)-5-(4-benzyl-1,3-thiazol-2-yl)-3-(pyrrolidin-2-ylmethyl)-1H-indole;
(R)-5-(3-benzyl-1,2,4-oxadiazol-5-yl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;
20 (R)-5-(3-benzyl-1,2,4-oxadiazol-5-yl)-3-(pyrrolidin-2-ylmethyl)-1H-indole;
(R)-5-(3-benzyl-1,2,4-oxadiazol-5-ylmethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole; and
(R)-5-(3-benzyl-1,2,4-oxadiazol-5-ylmethyl)-3-(pyrrolidin-2-ylmethyl)-1H-indole.

The present invention also relates to a pharmaceutical composition for treating
25 a condition selected from hypertension, depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain, and chronic paroxysmal hemicrania and headache associated with vascular disorders comprising an amount of a compound of the formula I or a pharmaceutically acceptable salt thereof effective in treating such condition and a pharmaceutically acceptable carrier.

30 The present invention also relates to a method for treating a condition selected from hypertension, depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain and chronic paroxysmal hemicrania and headache associated with vascular disorders comprising administering to a mammal (e.g., a human) requiring


-4-

such treatment an amount of a compound of the formula I or a pharmaceutically acceptable salt thereof effective in treating such condition.

The present invention also relates to a pharmaceutical composition for treating disorders arising from deficient serotonergic neurotransmission (e.g., depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain and chronic paroxysmal hemicrania and headache associated with vascular disorders) comprising administering to a mammal (e.g., a human) requiring such treatment an amount of a compound of the formula I or a pharmaceutically acceptable salt thereof effective in treating such condition.

The present invention also relates to a method for treating disorders arising from deficient serotonergic neurotransmission (e.g., depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain and chronic paroxysmal hemicrania and headache associated with vascular disorders) comprising administering to a mammal (e.g., a human) requiring such treatment an amount of a compound of the formula I or a pharmaceutically acceptable salt thereof effective in treating such condition.

The present invention also relates to a compound of the formula



25 where n , A , W , X , Y , Z , R_2 , R_3 , R_4 , R_5 , and R_{11} are as defined above; and R_{13} is C_1 - C_6 alkyl, aryl, or alkylaryl (preferably benzyl). When R_{11} is hydrogen, the epimers with the R absolute configuration at the chiral carbon site designated by an asterisk in formula II are preferred. When R_{11} is $-OR_{12}$ or $-NHCOR_{12}$ and n is 0 or 1, the epimers with the S absolute configuration at the chiral carbon site designated by an asterisk in formula


30 II are preferred. When R_{11} is $-OR_{12}$ or $-NHCOR_{12}$ and n is 2, the epimers with the R absolute configuration at the chiral carbon site designated by an asterisk in formula II are preferred. When R_{11} is $-OR_{12}$ or $-NHCOR_{12}$ and n is 0, the cis epimers [(2*S*, 3*S*) absolute configuration in the azetidine ring] are particularly preferred. When R_{11} is

-5-

-OR₁₂ or -NHCOR₁₂ and n is 1, the cis epimers [(2S, 4R) absolute configuration in the pyrrolidine ring] are particularly preferred. When R₁₁ is -OR₁₂ or -NHCOR₁₂ and n is 2, the cis epimers [(2R, 5R) absolute configuration in the piperidine ring] are particularly preferred. The compounds of formula II are useful as intermediates in preparing

5 compounds of formula I.

The present invention also relates to a compound of the formula



where n, A, W, X, Y, Z, R₂, R₃, R₄, R₅, R₁₁ and R₁₃ are as defined above; R₁₄ is halogen

15 (e.g. fluorine, chlorine, bromine, or iodine [preferably bromine or iodine]); and R₁₅ is -COCF₃, -SO₂CH₃, -SO₂Ph [Ph=phenyl], or -CO₂C(CH₃)₃ [preferably -COCF₃]. When R₁₁ is hydrogen, the epimers with the R absolute configuration at the chiral carbon site designated by an asterisk in formula III are preferred. When R₁₁ is -OR₁₂ or -NHCOR₁₂ and n is 0 or 1, the epimers with the S absolute configuration at the chiral carbon site


20 designated by an asterisk in formula III are preferred. When R₁₁ is -OR₁₂ or -NHCOR₁₂ and n is 2, the epimers with the R absolute configuration at the chiral carbon site designated by an asterisk in formula III are preferred. When R₁₁ is -OR₁₂ or -NHCOR₁₂ and n is 0, the cis epimers [(2S, 3S) absolute configuration in the azetidine ring] are particularly preferred. When R₁₁ is -OR₁₂ or -NHCOR₁₂ and n is 1, the cis epimers [(2S, 4R) absolute configuration in the pyrrolidine ring] are particularly preferred. When R₁₁

25 is -OR₁₂ or -NHCOR₁₂ and n is 2, the cis epimers [(2R, 5R) absolute configuration in the piperidine ring] are particularly preferred. The compounds of formula III are useful as intermediates in preparing compounds of formula II.

Detailed Description of the Invention


30 The compounds of the present invention are prepared via the following reaction scheme.

-6-



5


10



15

20

25




30


-7-

FORMULA II


IA
($R_1 = H$)



IB
($R_1 = CH_3$)



IC
(R_1 is not H)



5

10

15

Compounds of formula III can be prepared by the Mitsunobu coupling reaction of compounds of formulas IV and V wherein n, A, W, X, Y, Z, R₂, R₃, R₄, R₅, R₁₁, R₁₂, R₁₃, R₁₄, and R₁₅ are as defined above using a phosphine and an azodicarboxylate in a suitable solvent. Suitable phosphines include trialkyl phosphines and triarylphosphines, preferably triphenylphosphine. Suitable azodicarboxylates include dialkyl azodicarboxylates, preferably diethyl diazodicarboxylate. Suitable solvents include methylene chloride, ethers, (tetrahydrofuran, diethyl ether, and 1,4-dioxane), N,N-dimethylformamide and acetonitrile. The preferred solvent is tetrahydrofuran. The reaction is conducted at a temperature of from about 0°C to about 65°C, most preferably at about 25°C.

Compounds of formula II can be prepared by the transition metal catalyzed cyclization of compounds of the formula III where n, A, W, X, Y, Z, R₂, R₃, R₄, R₅, R₁₁, and R₁₃ are as defined above, R₁₄ is halogen (preferably bromine or iodine) and R₁₅ is -COCF₃, -SO₂CH₃, -SO₂Ph, or -CO₂C(CH₃)₃, preferably trifluoromethylacetyl [-COCF₃], in a suitable inert solvent with a phase transfer catalyst and a base. Suitable catalysts include palladium salts such as palladium (II) acetate or palladium (II) chloride (preferably palladium acetate) and rhodium salts, such as tris(triphenyl)rhodium (I) chloride. Suitable solvents include N,N-dimethylformamide, acetonitrile, and N-methylpyrrolidine. The preferred solvent is N,N-dimethylformamide. Suitable phase transfer catalysts include tetraalkylammonium halides, and tetra-n-butylammonium chloride preferably the latter. Suitable bases include tertiary amines, sodium hydrogen carbonate, and sodium carbonate. The preferred base is triethylamine. The reaction is conducted at a temperature of from about 60°C to about 180°C, preferably from about 80°C to 100°C.


Compounds of formula IB (R₁ = -CH₃) are prepared by hydride reduction of a compound of the formula II where n, A, W, X, Y, Z, R₂, R₃, R₄ and R₅ are as defined above, and R₁₃ is selected from C₁-C₆ alkyl, aryl, and alkylaryl (preferably benzyl) with a hydride reducing agent in an inert solvent. Suitable hydride reducing agents include lithium aluminum hydride, diborane, lithium borohydride, and sodium amide. The preferred reagent is lithium aluminum hydride. Suitable solvents include ethers, such as diethyl ether, tetrahydrofuran, 1,4-dioxane and 1,2-dimethoxyethane. The preferred solvent is tetrahydrofuran. The reduction is conducted at a temperature of from about 30°C to about 100°C, preferably from about 65°C to about 70°C.

Compounds of formula IA ($R_1=H$) are prepared by catalytic reduction of a compound of the formula II where n, A, W, X, Y, Z, R_2 , R_3 , R_4 , R_5 and R_{13} are as defined above under an atmosphere of hydrogen, preferably at a pressure of about 1 to about 3 atmospheres, or using a hydrogen source such as ammonium formate or formic acid in an inert solvent. Suitable catalysts include palladium on carbon, Raney nickel, and platinum oxide. The preferred catalyst is palladium on carbon. Suitable solvents include C_1 to C_6 alcohols, N,N-dimethylformamide, ethyl acetate, and acetonitrile. The preferred solvent is ethanol. The reaction is conducted at a temperature of from about 0°C to about 60°C, preferably about 25°C.

10 Compounds of formula IC ($R_1 \neq H$) are also prepared by the alkylation of a compound of the formula IA ($R_1=H$) wherein R_2 , R_3 , R_4 , R_5 , R_{11} , W, X, Y, Z, A, and n are as defined above with an alkylating agent of the formula R_1 -LG and a base in an inert solvent, where LG is a suitable leaving group and R_1 is as defined above except for hydrogen. Examples of suitable leaving groups include -I, -Br, -Cl, -OSO₂Ph, 15 -OSO₂CH₃, and -OSO₂CF₃. Suitable alkylating agents include alkyl halides (chlorides, bromides, or iodides), alkyl tosylates, alkyl mesylates, alkyl triflates, α,β -unsaturated ketones, α,β -unsaturated esters, α,β -unsaturated amides, and α,β -unsaturated nitriles. Alkyl halides (iodides) are preferred. Suitable solvents include methylene chloride, chloroform, carbon tetrachloride, acetonitrile, tetrahydrofuran, diethyl ether, dioxane, 20 N,N-dimethylformamide, ethanol, propanol, and methanol. The preferred solvent is acetonitrile. The reaction is conducted between a temperature of about 0°C to about 150°C, preferable about 25°C to about 65°C.

Compounds of formula V are prepared via the following reaction scheme.

-10-





5

10

15

20



Compounds of the formula VI can be prepared using the Wittig reaction in a suitable solvent involving compounds of the formulas VII and VIII wherein n and R_{13} are defined as above. R_{16} is C_1 - C_6 alkyl, aryl, or alkylaryl. Suitable solvents include ethers such as diethyl ether, tetrahydrofuran, and 1,4-dioxane. Tetrahydrofuran is the

preferred solvent. The reaction is conducted at a temperature of from about -78°C to about 30°C, preferably at about -78°C.

Compounds of the formula V can be prepared from a hydride reduction of a compound of formula VI wherein n, R₁₃ and R₁₆ are as defined above with a hydride
5 reducing agent in an inert solvent. Suitable hydride reducing agents include lithium aluminum hydride, lithium borohydride, sodium borohydride, and diisobutylaluminum hydride. The preferred reagent is diisobutylaluminum hydride. Suitable solvents include ethers, such as diethyl ether, tetrahydrofuran, 1,4-dioxane and 1,2-dimethoxyethane. The preferred solvent is tetrahydrofuran. The reduction is conducted
10 at a temperature of from about -100°C to about 0°C, preferably from about -80°C to about -70°C.


Compounds of the formula VII can be prepared using methods known to one skilled in the art, such as, for example, as outlined in S. Kiyooka, et al., J. Org. Chem., 5409 (1989) and Y. Hamada, et al., Chem. Pharm. Bull., 1921 (1982).

15 Compounds of the formula VIII are either commercially available or can be prepared using methods known to one skilled in the art, such as, for example, as outlined in L. Fieser and M. Fieser, Reagents for Organic Synthesis, John Wiley and Sons, New York, Vol. 1, p. 112 (1967).

Compounds of formula IV are prepared using the following reaction scheme.


-12-

5



X I


10



I X

15

20



I V

25

Compounds of formula IX can be prepared by reacting a compound of formula XI wherein A, W, X, Y, Z, R₂, R₃, R₄ and R₅ are as defined above with either chlorine, bromine, or iodine in a suitable solvent with a suitable base. Reaction with bromine is preferred. Suitable solvents include C₁ - C₆ alcohols, methylene chloride, chloroform, or carbon tetrachloride. The preferred solvent is methanol. Suitable bases include triethylamine, pyridine, sodium carbonate, and sodium hydrogen carbonate. The preferred base is sodium hydrogen carbonate. The reaction is conducted at a temperature of from about 0°C to about 65°C, preferably at about 25°C.

-13-

Compounds of formula IV can be prepared by reacting a compound of formula IX wherein A, W, X, Y, Z, R₂, R₃, R₄, R₅, and R₁₄ are as defined above with the acid chloride or symmetrical anhydride of the formula R₁₅CO₂H in a suitable solvent with an suitable base. The preferred acid chloride or anhydride is trifluoroacetic anhydride.

5 Suitable solvents include methylene chloride, chloroform as well as ethers, including tetrahydrofuran, diethyl ether and 1,4-dioxane. The preferred solvent is methylene chloride. Suitable bases include triethylamine, pyridine, and sodium hydrogen carbonate. The preferred base is pyridine. The reaction is conducted at a temperature of from about 0°C to about 65°C, preferably at about 25°C.

10 Compounds of the formula XI can be prepared using methods known to one skilled in the art, such as, for example, as outlined in European Patent Application Pub. No. 0 438 230 A2.

Compounds of formula IX where W is oxygen, X and Z are nitrogen, and Y is carbon can also be prepared by reacting together compounds of the formula




20 wherein A, R₄, R₁₂ are as defined above, and R₁₇ is C₁-C₈ alkyl or aryl in an inert solvent in the presence of a base [P. Sauerberg, et al. J. Med. Chem., 687 (1991), G.A. Showell, J. Med. Chem., 1086 (1991) and European Patent Application Pub. No. 0 438 230 A2]. Suitable solvents include ethers such as tetrahydrofuran, 1,4-dioxane, and diethyl ether, methylene chloride, chloroform, carbon tetrachloride, and C₁-C₈ alcohols.

25 The preferred solvent is tetrahydrofuran. Suitable bases included sodium metal, sodium hydride, potassium hydride, and potassium *t*-butoxide. The preferred base is sodium hydride. The reaction is conducted at a temperature of about 0°C to 101°C, preferably at about 66°C.

30 Compounds of formula XII, if not commercially available, can be prepared via the reaction of a compound of the formula

-14-



5

wherein A and R₁₇ are as defined above with either chlorine, bromine, or iodine in a suitable solvent with a suitable base. Reaction with bromine is preferred. Suitable solvents include C₁-C₆ alcohols, methylene chloride, chloroform, or carbon tetrachloride. The preferred solvent is methanol. Suitable bases include triethylamine, pyridine, sodium carbonate, and sodium hydrogen carbonate. The preferred base is sodium hydrogen carbonate. The reaction is conducted at a temperature of about 0°C to about 65°C, most preferably at about 25°C.

Compounds of the formula XIII can be prepared using methods known to one skilled in the art, such as, for example, C.L. Bell, et al. J. Org. Chem., 2873 (1964).

Compounds of formula XIV are available either commercially or using methods known to one skilled in the art, such as, for example, E. Ferber, et al., Chem. Ber., 839 (1939).

The compounds of the formula I which are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially isolate a compound of the formula I from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent, and subsequently convert the free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is obtained.

The acids which are used to prepare the pharmaceutically acceptable acid addition salts of the base compounds of this invention are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate or bisulfate, phosphate or

acid phosphate, acetate, lactate, citrate or acid citrate, tartrate or bitartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)] salts.

Those compounds of the formula I which are also acidic in nature, i.e., where
5 R_2 contains a carboxylate, are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline-earth metal salts and particular, the sodium and potassium salts. These salts are all prepared by conventional techniques. The chemical bases which are used as reagents to prepare the pharmaceutically acceptable base salts of this invention are
10 those which form non-toxic base salts with the herein described acidic compounds of formula I. These non-toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium, calcium, magnesium, etc. These salts can easily be prepared by treating the corresponding acidic compounds with an aqueous solution containing the desired pharmacologically acceptable cations,
15 and then evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, they may also be prepared by mixing lower alkanolic solutions of the acidic compounds and the desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are preferably employed in order to ensure
20 completeness of reaction of maximum product of yields of the desired final product.

The compounds of the formula I and the pharmaceutically acceptable salts thereof (hereinafter, also referred to as the active compounds of the invention) are useful psychotherapeutics and are potent serotonin ($5-HT_1$) agonists and may be used in the treatment of depression, anxiety, eating disorders, obesity, drug abuse, cluster
25 headache, migraine, chronic paroxysmal hemicrania and headache associated with vascular disorders, pain, and other disorders arising from deficient serotonergic neurotransmission. The compounds can also be used as centrally acting antihypertensives and vasodilators. The active compounds of the invention can be evaluated as anti-migraine agents by testing the extent to which they mimic sumatriptan
30 in contracting the dog isolated saphenous vein strip [P.P.A. Humphrey et al., Br. J. Pharmacol., 94, 1128 (1988)]. This effect can be blocked by methiothepin, a known serotonin antagonist. Sumatriptan is known to be useful in the treatment of migraine and produces a selective increase in carotid vascular resistance in the anesthetized

-16-

dog. It has been suggested [W. Fenwick et al., Br. J. Pharmacol., 96, 83 (1989)] that this is the basis of its efficacy.

The serotonin 5-HT₁ agonist activity can be measured in in vitro receptor binding assays as described for the 5-HT_{1A} receptor using rat cortex as the receptor source and [3H]-8-OH-DPAT as the radioligand [D. Hoyer et al. Eur. J. Pharm., Vol. 118, 13 (1985)] and as described for the 5-HT_{1D} receptor using bovine caudate as the receptor source and [3H]serotonin as the radioligand [R.E. Heuring and S.J. Peroutka, J. Neuroscience, Vol. 7, 894 (1987)] 5-HT₁ agonist activity is defined by agents with affinities (IC₅₀'s) of 250nM or less at either binding assay.

10 The compositions of the present invention may be formulated in a conventional manner using one or more pharmaceutically acceptable carriers. Thus, the active compounds of the invention may be formulated for oral, buccal, intranasal, parenteral (e.g., intravenous, intramuscular or subcutaneous) or rectal administration or in a form suitable for administration by inhalation or insufflation.

15 For oral administration, the pharmaceutical compositions may take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g. pregelatinized maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g. lactose, microcrystalline cellulose or calcium phosphate); lubricants (e.g. magnesium stearate, 20 talc or silica); disintegrants (e.g. potato starch or sodium starch glycolate); or wetting agents (e.g. sodium lauryl sulphate). The tablets may be coated by methods well known in the art. Liquid preparations for oral administration may take the form of, for example, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations 25 may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g. sorbitol syrup, methyl cellulose or hydrogenated edible fats); emulsifying agents (e.g. lecithin or acacia); non-aqueous vehicles (e.g. almond oil, oily esters or ethyl alcohol); and preservatives (e.g. methyl or propyl p-hydroxybenzoates or sorbic acid).

30 For buccal administration the composition may take the form of tablets or lozenges formulated in conventional manner.

The active compounds of the invention may be formulated for parenteral administration by injection, including using conventional catheterization techniques or

infusion. Formulations for injection may be presented in unit dosage form e.g. in ampules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulating agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for reconstitution with a suitable vehicle, e.g. sterile pyrogen-free water, before use.

The active compounds of the invention may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

For intranasal administration or administration by inhalation, the active compounds of the invention are conveniently delivered in the form of a solution or suspension from a pump spray container that is squeezed or pumped by the patient or as an aerosol spray presentation from a pressurized container or a nebulizer, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. The pressurized container or nebulizer may contain a solution or suspension of the active compound. Capsules and cartridges (made, for example, from gelatin) for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

A proposed dose of the active compounds of the invention for oral, parenteral or buccal administration to the average adult human for the treatment of the conditions referred to above (e.g., migraine) is 0.1 to 200 mg of the active ingredient per unit dose which could be administered, for example, 1 to 4 times per day.

Aerosol formulations for treatment of the conditions referred to above (e.g., migraine) in the average adult are preferably arranged so that each metered dose or "puff" of aerosol contains 20 μ g to 1000 μ g of the compound of the invention. The overall daily dose with an aerosol will be within the range 100 μ g to 10 mg. Administration may be several times daily, for example 2, 3, 4 or 8 times, giving for example, 1, 2 or 3 doses each time.

The following Examples illustrate the preparation of the compounds of the present invention. Melting points are uncorrected. NMR data are reported in parts per million (d) and are referenced to the deuterium lock signal from the sample solvent.

Specific rotations were measured at room temperature using the sodium D line (589 nm). Unless otherwise stated, all mass spectrum were performed using electron impact (EI, 70 eV) conditions.

Commercial reagents were utilized without further purification. Chromatography
5 refers to column chromatography performed using 32-63 μ m silica gel and executed under nitrogen pressure (flash chromatography) conditions. Room temperature refers to 20-25°C.

Example 1

General Procedure for the Hydride Reduction of 3-(N-Benzyloxycarbonyl- 10 pyrrolidin-2-ylmethyl)-1H-indoles Forming 3-(N-Methylpyrrolidin-2-ylmethyl)- 1H-indoles

To a stirred mixture of lithium aluminum hydride (0.152 g, 4.00 mmol, 2 eq.) in anhydrous tetrahydrofuran (10 mL) at 0°C was added rapidly a solution of the 3-(N-benzyloxycarbonyl-pyrrolidin-1-ylmethyl)-1H-indole (2.00 mmol) in anhydrous
15 tetrahydrofuran (5 mL). The resulting mixture is heated at reflux under a nitrogen atmosphere for 3 hours. The reaction mixture is cooled, and water (0.25 mL), 15% aqueous sodium hydroxide (0.25 mL), and then more water (0.75 mL) were added sequentially. The resulting mixture was stirred at 25°C for 30 minutes, filtered, and the filtrate was evaporated under reduced pressure. The residue was column
20 chromatographed using silica gel (approximately 50 g) and elution with a solution methylene chloride: methanol: ammonium hydroxide [9: 1: 0.1] or other appropriate solvent system to afford the corresponding 3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole.

Following this procedure the following compounds were prepared:

A. (R)-5-(4-benzyl-1,3-thiazol-2-yl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole
25 (R)-5-(4-Benzyl-1,3-thiazol-2-yl)-3-(N-benzyloxycarbonylpyrrolidin-2-yl-methyl)-1H-
indole was used. Chromatography using 5% triethylamine in ethyl acetate afforded the title compound (71%) as a white solid: mp, 146.0-148.0°C; ¹³C NMR (CDCl₃) δ 169.8, 157.1, 139.3, 137.3, 129.2, 128.5, 128.0, 126.4, 125.7, 123.2, 121.2, 117.6, 114.8, 113.2, 111.5, 66.6, 57.5, 40.8, 38.1, 31.4, 29.6, 21.9; LRMS (m/z, relative intensity) 387 (M⁺, 4),
30 303 (34), 155 (30), 147 (17), 115 (18), 85 (63), 84 (100), 83 (57); [α]²⁵ = +68° (CHCl₃, c=1.0). Anal. calcd. for C₂₄H₂₅N₃S • 1/4 H₂O: C, 73.54; H, 6.56; N, 10.72. Found: C, 73.50; H, 6.53; N, 10.57.

B. (R)-5-(3-benzyl-1,2,4-oxadiazol-5-yl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole

(R)-5-(3-Benzyl-1,2,4-oxadiazol-5-yl)-3-(N-benzyloxycarbonylpyrrolidin-2-ylmethyl)-1H-indole was used. Column chromatography as described above afforded the title compound (34%) as a tan solid: ¹H NMR (CDCl₃) δ 8.48 (s, 1H), 8.36 (s, 1H), 7.91 (dd, J=8 and 2 Hz, 1H), 7.43-7.25 (m, 6H), 7.12 (s, 1H), 4.13 (s, 2H), 3.28-3.15 (m, 2H), 2.77-2.68 (m, 1H), 2.53 (m, 1H), 2.46 (s, 3H), 2.26 (q, J=8 Hz, 1H), 1.92-1.74 (m, 2H), 1.74-1.54 (m, 2H); HRMS, calculated for C₂₃H₂₄N₄O 372.1945, found 372.1978.

C. (R)-5-(3-benzyl-1,2,4-oxadiazol-5-ylmethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole

(R)-5-(3-Benzyl-1,2,4-oxadiazol-5-ylmethyl)-3-(N-benzyloxycarbonylpyrrolidin-2-ylmethyl)-1H-indole was used. Column chromatography as described above afforded the title compound (24%) as a beige resin: ¹H NMR (CDCl₃) δ 8.10 (br s, 1H), 7.47 (s, 1H), 7.34-7.18 (m, 6H), 7.08 (dd, J=8 and 2 Hz, 1H), 7.04 (br s, 1H), 4.25 (s, 2H), 4.01 (s, 2H), 3.22-3.07 (m, 2H), 2.66-2.55 (m, 1H), 2.54-2.43 (m, 1H), 2.42 (s, 3H), 2.24 (q, J=8 Hz, 1H), 1.86-1.69 (m, 2H), 1.68-1.50 (m, 2H); HRMS calculated for C₂₄H₂₆N₄O 386.2070, found 386.2074.

Example 2

General Procedure for the Catalytic Reduction of 3-(N-Benzyloxycarbonylpyrrolidin-2-ylmethyl)-1H-indoles Forming 3-(Pyrrolidin-2-ylmethyl)-1H-indoles

A mixture of the 3-(N-benzyloxycarbonylpyrrolidin-2-ylmethyl)-1H-indole (2.00 mmol) and 10% palladium on carbon (0.20 g) in absolute ethanol (15 mL) was shaken under a hydrogen atmosphere (3 atm) for 4-24 hours, depending on substrate. The resulting reaction mixture was filtered through diatomaceous earth, and the filtrate was evaporated under reduced pressure. The residue was column chromatographed using silica gel (approximately 10 g) and elution with a solution of methylene chloride: methanol: ammonium hydroxide [8: 2: 0.2] or other appropriate solvent system to afford the corresponding 3-(pyrrolidin-2-ylmethyl)-1H-indole.

Following this procedure the following compounds were prepared:

A. (R)-5-(4-benzyl-1,3-thiazol-2-yl)-3-(pyrrolidin-2-ylmethyl)-1H-indole

(R)-5-(4-Benzyl-1,3-thiazol-2-yl)-3-(N-benzyloxycarbonylpyrrolidin-2-yl-methyl)-1H-indole was used, and the reaction was heated at 40°C for 24 hours. Chromatography using methylene chloride: methanol: ammonium hydroxide [9: 1: 0.1] afforded the title

-20-

compound (12%) as an amorphous solid: ^1H NMR (CDCl_3) δ 9.1 (br s, indole NH), 8.17 (d, $J=1.4$ Hz, 1H), 7.74 (dd, $J=1.6$ and 8.5 Hz, 1H), 7.35-7.21 (m, 6H), 7.02 (s, 1H), 6.67 (s, 1H), 4.22 (s, 2H), 3.5 (br s, NH), 3.41-3.29 (m, 1H), 3.03-2.73 (m, 4H), 1.94-1.61 (m, 3H), 1.49-1.38 (m, 1H); ^{13}C NMR (CDCl_3) δ 169.9, 157.0, 139.2, 137.4, 129.2, 128.5, 127.7, 126.4, 125.5, 123.8, 121.2, 117.3, 114.3, 113.3, 111.7, 59.2, 46.0, 38.1, 31.5, 31.1, 24.9; HRMS calculated for $\text{C}_{23}\text{H}_{23}\text{N}_3\text{S}$ 374.1615, found 374.1691.

Example 3

General Procedure for the Formation of 3-(N-Benzylloxycarbonylpyrrolidin-2-ylmethyl)-1H-indoles Via the Palladium Catalyzed Cyclization of 1-(N-Benzylloxycarbonylpyrrolidin-2-yl)-3-(N-(2-halophenyl)-N-trifluoroacetyl-amino)-propenes

A mixture of the 1-(N-benzylloxycarbonylpyrrolidin-2-yl)-3-(N-(2-halophenyl)-N-trifluoroacetyl-amino)propene (2.00 mmol), tetrabutylammonium chloride (2.00 mmol), and palladium(II) acetate (0.089 g, 0.40 mmol, 0.2 eq) in a solution of triethylamine (8 mL) and anhydrous N,N-dimethylformamide (4 mL) was heated at reflux under nitrogen for 2 hours. The resulting reaction mixture was evaporated under reduced pressure, and the residue was partitioned between ethyl acetate (25 mL) and water (25 mL). The ethyl acetate layer was removed, and the aqueous layer was extracted with ethyl acetate (25 mL). The organic extracts were combined, dried (MgSO_4), and evaporated under reduced pressure. The residue was column chromatographed using silica gel (approximately 50 g) and elution with 40% ethyl acetate in hexanes or an appropriate solvent system to afford the corresponding 3-(N-benzylloxycarbonylpyrrolidin-2-ylmethyl)-1H-indole.

Following this procedure the following compounds were prepared:

A. (R)-5-(4-Benzyl-1,3-thiazol-2-yl)-3-(N-benzylloxycarbonylpyrrolidin-2-ylmethyl)-1H indole

(R)-1-(N-Benzylloxycarbonylpyrrolidin-2-yl)-3-(N-(2-bromo-4-(4-benzyl-1,3-thiazol-2-yl)phenyl)-N-trifluoroacetyl-amino)propene was used. Chromatography using elution with ethyl acetate gradient in hexanes [1:3 to 2:5] afforded the title compound (58%) as a pale yellow foam: FAB LRMS (m/z , relative intensity) 509 (MH^+ , 37), 508 (M^+ , 100), 462 (5), 372 (8), 304 (33); FAB HRMS calculated for $[\text{C}_{31}\text{H}_{30}\text{N}_3\text{O}_2\text{S}\cdot\text{H}]^+$ 509.2139, found 509.2106. Anal. calcd for $\text{C}_{31}\text{H}_{30}\text{N}_3\text{O}_2\text{S} \cdot 1/2 \text{C}_4\text{H}_8\text{O}_2$ [ethyl acetate]: C, 71.71; H, 6.20; N, 7.60. Found: C, 71.55; H, 5.82; N, 7.64.

B. (R)-5-(3-Benzyl-1,2,4-oxadiazol-5-yl)-3-(N-benzyloxycarbonylpyrrolidin-2-ylmethyl)-1H-indole

(R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-bromo-4-(3-benzyl-1,2,4-oxadiazol-5-yl)phenyl)-N-trifluoroacetyl-amino)propene was used. Column chromatography using the above described solvent system afforded the title compound (74%) as a light yellow resin: $R_f = 0.41$ (hexane-EtOAc 50:50); HRMS calculated for $C_{30}H_{29}N_4O_3$ 493.2288, found: 493.2240.

C. (R)-5-(3-Benzyl-1,2,4-oxadiazol-5-ylmethyl)-3-(N-benzyloxycarbonylpyrrolidin-2-yl-methyl)-1H-indole

(R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-bromo-4-(3-benzyl-1,2,4-oxadiazol-5-ylmethyl)phenyl)-N-trifluoroacetyl-amino)propene was used. Column chromatography using the above described solvent system afforded the title compound (61%) as a tan resin: $R_f = 0.063$ (hexane-EtOAc 50:50); HRMS calculated for $C_{31}H_{31}N_4O_3$ 507.2396, found: 507.2387.

Example 4

General Procedure for the Formation 1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-2-halophenyl)-N-trifluoroacetyl-amino)propenes from the Mitsunobu Coupling of 2-Halo-N-trifluoroacetylanilines with 1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene.

To a stirred solution of 1-(N-benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene (R, or S, or racemate, 2.00 mmol), the 2-halo-N-trifluoroacetylaniline (2.5 mmol, 1.25 eq), and triphenylphosphine (0.655 g, 2.50 mmol, 1.25 eq) in anhydrous tetrahydrofuran (15 mL) at 0°C under a nitrogen atmosphere was added diethyl azodicarboxylate (0.39 mL, 2.48 mmol, 1.25 eq) dropwise. The reaction solution was slowly warmed to 25°C over the course of 2 hours, and then stirred at 25°C under a nitrogen atmosphere for an additional 12 hours. The resulting reaction solution was evaporated under reduced pressure, and the residue was column chromatographed using silica gel (approximately 150 g) and elution with an appropriate solvent system to afford the corresponding 1-(N-benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-halophenyl)-N-trifluoroacetyl-amino)propene.

Following this procedure the following compounds were prepared:

A. (R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-bromo-4-(4-benzyl-1,3-thiazol-2-yl)phenyl)-N-trifluoroacetyl-amino)propene

4-(4-Benzyl-1,3-thiazol-2-yl)-2-bromo-1-trifluoroacetylaminobenzene and (R)-1-(N-benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene were used. Chroma-

-22-

tography using elution with a 1-5% ether gradient in methylene chloride afforded the title compound (97%) as a white foam: FAB LRMS (m/z , relative intensity) 686 (MH_2^+ , 100), 685 (MH^+ , 60), 684 (M^+ , 90), 640 (23), 578 (15), 441 (17), 371 (20); FAB HRMS calculated for $[C_{33}H_{29}BrF_3N_3O_3S \cdot H]^+$ [with ^{79}Br and ^{32}S] 684.1145, found 684.1157.

5 B. (R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-bromo-4-(3-benzyl-1,2,4-oxadiazol-5-yl))phenyl)-N-trifluoroacetylaminopropene

4-(3-Benzyl-1,2,4-oxadiazol-5-yl)-2-bromo-1-trifluoroacetylaminobenzene and (R)-1-(N-benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene were used. Chromatography using elution with 5% ether in methylene chloride afforded the title
10 compound (88%) as a thick yellow oil: R_f = 0.32 ($CHCl_3$); LRMS (m/z , relative intensity) 669 (M^+ , 25).

 C. (R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-(N-(2-bromo-4-(3-benzyl-1,2,4-oxadiazol-5-ylmethyl))phenyl)-N-trifluoroacetylaminopropene

4-(3-Benzyl-1,2,4-oxadiazol-5-ylmethyl)-2-bromo-1-trifluoroacetylaminobenzene
15 and (R)-1-(N-benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene were used. Chromatography using elution with 5% ether in methylene chloride afforded the title compound (90%) as a thick yellow oil: R_f = 0.75 ($CHCl_3$ - CH_3OH 20:1); LRMS (m/z , relative intensity) 683 (M^+ , 18).

Example 5

20 (R)-1-(N-Benzyloxycarbonylpyrrolidin-2-yl)-3-hydroxypropene

To a stirred solution of ethyl (R)-3-(N-benzyloxycarbonylpyrrolidin-2-yl)-2-propenoate (3.03 g, 10.00 mmol) in anhydrous tetrahydrofuran (75 mL) at $-78^\circ C$ under nitrogen was added dropwise a solution of diisobutylaluminum hydride (1.0 M in hexanes, 22.0 mL, 22.0 mmol, 2.2 eq). The resulting solution was stirred at $-78^\circ C$
25 under nitrogen for 30 minutes. The reaction solution was then allowed to warm to room temperature over the course of 2 hours. A saturated solution of sodium hydrogen carbonate (50 mL) was added, and the aqueous mixture was extracted with ethyl acetate (3 x 50 mL). The extracts were combined, dried ($MgSO_4$), and evaporated under reduced pressure. Column chromatography of the residue with an diethyl
30 ether/hexanes [1:1] afforded the title compound as a clear colorless oil (1.41 g, 5.40 mmol, 54%): 1H NMR ($CDCl_3$) δ 7.40-7.25 (m, 5H), 5.75-5.53 (m, 2H), 5.20-5.00 (m, 2H), 4.38 (br m, 1H), 4.06 (br d, J =13.7 Hz, 3H), 3.45 (br t, J =7.0 Hz, 1H), 2.03-1.68

-23-

(m, 4H); $[\alpha]^{25} = +34^\circ$ (MeOH, c=1.0); HRMS calculated for $C_{15}H_{19}NO_3$ 261.1365, found 261.1356.

Example 6

Ethyl (R)-3-(N-Benzoyloxycarbonylpyrrolidin-2-yl)-2-propenoate

5 To a stirred solution of N-carbobenzoyloxypyrrolidine-2-carboxaldehyde (1.17 g, 5.00 mmol) in anhydrous tetrahydrofuran at -78°C was added (carboethoxymethylene)-triphenylphosphorane (2.09 g, 6.00 mmol, 1.2 eq) as a solid portionwise. The resulting reaction mixture was stirred at room temperature under nitrogen for 2 hours, and then heated at reflux under nitrogen for 1 hour. The reaction mixture was evaporated under
10 reduced pressure and the residue was column chromatographed using silica gel (approximately 100 g) and elution with 20% diethyl ether in hexanes to afford the title compound as a clear, colorless oil (1.11 g, 3.65 mmol, 73%): ^1H NMR (CDCl_3) δ 7.34-7.25 (m, 5H), 6.89-6.76 (m, 1H), 5.88-5.74 (m, 1H), 5.18-5.05 (m, 2H), 4.60-4.43 (m, 1H), 4.17 (q, $J=7.1$ Hz, 2H), 3.55-3.40 (m, 2H), 2.11-2.00 (m, 1H), 1.90-1.75 (m, 3H), 1.28
15 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (CDCl_3) [Note: due to slow nitrogen inversion two conformers of the products are seen by NMR spectroscopy] δ 166.3, 154.7, 147.9, 147.4, 136.6, 128.4, 127.9, 120.9, 66.9, 65.8, 60.4, 58.1, 57.7, 46.8, 46.4, 31.6, 30.8, 23.6, 22.8, 22.6, 15.3, 14.2.

Example 7

General Synthesis of 2-Halo-N-trifluoroacetylanilines from Reaction of 2-Haloanilines and Trifluoroacetic Anhydride

20 To a stirred solution of the 2-haloaniline (2.00 mmol) and pyridine (0.18 mL, 2.22 mmol, 1.1 eq) in anhydrous methylene chloride (10 mL) at 0°C under a nitrogen atmosphere was added dropwise trifluoroacetic anhydride (0.31 mL, 2.19 mmol, 1.1
25 eq). The resultant reaction mixture was stirred at 0°C under a nitrogen atmosphere for 3 hours. A saturated solution of sodium hydrogen carbonate was added (15 mL), and this aqueous mixture was extracted with ethyl acetate (3 x 15 mL). The extracts were combined, dried (MgSO_4), and evaporated under reduced pressure. If necessary, the residue was column chromatographed using silica gel (approximately 50 g) and elution
30 with an ethyl acetate gradient in hexanes to afford the corresponding 2-halo-N-trifluoroacetylaniline.

-24-

Following this procedure the following compounds were prepared:

A. 4-(4-Benzyl-1,3-thiazol-1-yl)-2-bromo-1-trifluoroacetylaminobenzene

4-(4-Benzyl-1,3-thiazol-2-yl)-2-bromoaniline was used. The extraction residue was triturated in diethyl ether/hexanes [1:1, 10 mL] to afford the title compound (92%)
5 as a white powder: mp, 102.0-104.0°C; ¹³C NMR (CDCl₃) δ 164.9, 158.0, 138.7, 134.1, 132.6, 130.1, 129.1, 128.6, 126.8, 126.6, 121.8, 115.2, 114.4, 38.0. Anal. calcd for C₁₈H₁₂F₃BrN₂OS: C, 48.99; H, 2.74; N, 6.35. Found: 48.72; H, 2.58; N, 6.29.

B. 4-(3-Benzyl-1,2,4-oxadiazol-5-yl)-2-bromo-1-trifluoroacetylaminobenzene

4-(3-Benzyl-1,2,4-oxadiazol-5-yl)-2-bromoaniline was used. Column
10 chromatography as described above afforded the title compound (81%) as a white solid: mp, 152.0-153.0°C; ¹H NMR (CDCl₃) δ 8.64 (br s, 1H), 8.53 (d, J=8 Hz, 1H), 8.38 (d, J=2 Hz, 1H), 8.13 (dd, J=8 and 2 Hz, 1H), 7.40-7.26 (m, 5H), 4.14 (s, 2H); LRMS (m/z, relative intensity) 426 (M+,85).

C. 4-(3-Benzyl-1,2,4-oxadiazol-5-ylmethyl)-2-bromo-1-trifluoroacetylaminobenzene
15 benzene

4-(3-Benzyl-1,2,4-oxadiazol-5-ylmethyl)-2-bromoaniline was used. Column chromatography as described above afforded the title compound (90%) as a yellow resin: ¹H NMR (CDCl₃) δ 8.59 (br s, 1H); 8.36 (br s, 1H), 8.22 (d, J=8 Hz, 1H), 7.42 (s, 1H), 7.24-7.32 (m, 5H), 4.10 (s, 2H), 4.01 (s, 2H); LRMS (m/z, relative intensity) 440
20 (M+,90).

Example 8

4-(4-Benzyl-1,3-thiazol-2-yl)-2-bromoaniline

A mixture of 4-amino-3-bromobenzthioamide (1.66 g, 7.18 mmol) and 1-chloro-3-phenylacetone [Tarhouni, R. et al., Tetrahedron Letters, 835 (1984)] (1.36 g, 8.07 mmol,
25 1.1 eq) in absolute ethanol (18 mL) was heated at reflux under nitrogen for 2.5 hours. The resulting reaction mixture was evaporated under reduced pressure, and the residue was partitioned between ethyl acetate (20 mL) and a saturated solution of sodium hydrogen carbonate (20 mL). The ethyl acetate layer was removed, and the aqueous layer was extracted with ethyl acetate (2 x 20 mL). The organic extracts were
30 combined, dried (MgSO₄), and evaporated under reduced pressure. The residual solid was chromatographed using silica gel (approximately 175 g) and elution with an ethyl acetate gradient in hexanes [1:4 to 1:1] to afford the title compound (68%) as a pale yellow solid: mp, 110-115°C; ¹³C NMR (CDCl₃) δ 166.8, 157.1, 145.6, 139.1, 130.7,

-25-

129.1, 128.6, 126.9, 126.4, 125.4, 115.3, 113.2, 109.2, 38.0. Anal. calcd for $C_{18}H_{13}BrN_2S$: C, 55.66; H, 3.79; N, 8.11. Found: C, 55.36; H, 3.71; N, 7.92.

Example 9

4-Amino-3-Bromobenzthioamide

5 A stirred solution of 4-amino-3-bromobenzonitrile (6.92 g, 35.1 mmol) and diethyl dithiophosphate (17.7 mL, 105 mmol, 3 eq.) in absolute ethanol (160 mL) at 0°C was perfused with hydrogen chloride gas at a moderate rate for 30 minutes. The resulting reaction mixture was stirred at room temperature for 12 hours, and then solvent was removed via evaporation under reduced pressure. The residue was suspended in a
10 saturated solution of sodium hydrogen carbonate (25 mL), and this aqueous mixture was extracted with ethyl acetate (3 x 25 mL). The organic extracts were combined, dried ($MgSO_4$), and evaporated under reduced pressure. The residue was chromatographed using silica gel (approximately 300 g) and elution with an acetone gradient in methylene chloride [1:50 to 1:20] to afford the title compound (1.02 g, 25%)
15 as an amorphous yellow solid: 1H NMR ($DMSO-d_6$) δ 9.41 (br s, NH), 9.13 (br s, NH), 8.11 (d, $J=2.1$ Hz, 1H), 7.78 (dd, $J=2.1$ and 8.6 Hz, 1H), 6.72 (d, $J=8.7$ Hz, 1H), 6.03 (s, 2NH); TLC: $R_F=0.15$ [1% diethyl ether in methylene chloride].

Example 10

General Procedure for the Formation of 2-Halo-4-(1,2,4-oxadiazol-5-yl)anilines
20 or 2-Halo-4-(1,2,4-oxadiazol-5-ylmethyl)anilines from the Condensation of the
Corresponding Alkyl 4-Amino-3-halobenzoates or Alkyl 2-(4-Amino-3-
halophenyl)acetates, respectively, with Phenylacetamide Oxime

Sodium hydride (87 mg of an 60% oil dispersion, 2 mmol) was added to a stirred solution of phenylacetamide oxime (0.33 g, 2.2 mmol, 1.1 eq) [C. L. Bell, et al.,
25 J. Org. Chem., 2873 (1964)] in anhydrous THF (10 ml), and the resulting reaction mixture was heated at reflux for 30 minutes. A solution of the alkyl 4-amino-3-halobenzoate or alkyl 2-(4-amino-3-halophenyl)acetate (1 mmol) in anhydrous THF (5 mL) was then added, and the reaction was heated under reflux for 2 hours. The mixture was allowed to cool to room temperature before water (10 ml) was added. The
30 resulting aqueous mixture was extracted with dichloromethane (3 x 25 ml). The extracts were combined, dried ($MgSO_4$), and evaporated under reduced pressure. The residue was chromatographed using silica gel (20 g) and elution with chloroform to afford the

corresponding 2-halo-4-(1,2,4-oxadiazol-5-yl)aniline or 2-halo-4-(1,2,4-oxadiazol-5-ylmethyl)aniline, respectively.

Following this procedure the following compounds were prepared:

A. 4-(3-Benzyl-1,2,4-oxadiazol-5-yl)-2-bromoaniline

5 Methyl 4-amino-3-bromobenzoate was used. Column chromatography as described above afforded the title compound (33%) as a tan solid; mp 144-145°C; ¹H NMR (CDCl₃) δ 8.18 (d, *J*=2 Hz, 1H), 7.82 (dd, *J*=8 and 2 Hz, 1H), 7.39-7.25 (m, 5H), 6.77 (d, *J*=8 Hz, 1H), 4.09 (s, 2H); LRMS (m/z, relative intensity) 330 (M+,90).

B. 4-(3-Benzyl-1,2,4-oxadiazol-5-ylmethyl)-2-bromoaniline

10 Ethyl 2-(4-amino-3-bromophenyl)acetate was used. Column chromatography as described above afforded the title compound (41%) as a yellow resin; ¹H NMR (CDCl₃) δ 7.34-7.24 (m, 6H), 7.00 (dd, *J*=8 and 2 Hz, 1H), 6.69 (d, *J*=8 Hz, 1H), 4.02 (s, 2H), 4.01 (s, 2H); LRMS (m/z, relative intensity) 334 (M+,15).

Example 11

15 General Procedure for the Bromination of Anilines to Form 2-Bromoanilines

To a stirred mixture of the aniline (2.00 mmol) and sodium hydrogen carbonate (0.21 g, 2.50 mmol, 1.25 eq) in methanol (10 mL) at 0°C was added dropwise bromine (0.113 mL, 2.19 mmol, 1.1 eq). The resulting reaction mixture was then stirred at 25°C for 30 minutes. The reaction mixture was then evaporated under reduced pressure, and
20 the residue was placed in a saturated solution of sodium hydrogen carbonate (10 mL). This aqueous mixture was extracted with ethyl acetate (3 x 15 mL). The extracts were combined, dried (MgSO₄), and evaporated under reduced pressure. The residue was column chromatographed using silica gel (approximately 50 g) and elution with an appropriate solvent system to afford the corresponding 2-bromoaniline.

25 Following this procedure the following compounds were prepared:

A. 4-Amino-3-bromobenzonitrile

4-Aminobenzonitrile was used. Chromatography using elution with a gradient of ethyl acetate in hexanes [1:5 to 1:3] afforded the title compound (71%) as a white solid: ¹H NMR (CDCl₃) δ 7.65 (d, *J*=2.1 Hz, 1H), 7.34 (dd, *J*=2.1 and 8.1 Hz, 1H), 6.71
30 (d, *J*=8.0 Hz, 1H), 4.6(br s, 2NH); TLC:R_f=0.25 [ethyl acetate/hexanes, 1:3].

B. Methyl 4-amino-3-bromobenzoate

Methyl 4-aminobenzoate was used. Chromatography using elution with ethyl acetate in hexanes [1:4] afforded the title compound (36%) as an orange oil: ¹H NMR

-27-

(CDCl₃) δ 8.09 (d, $J=2$ Hz, 1H), 7.75 (dd, $J=9$ and 2 Hz, 1H), 6.71 (d, $J=9$ Hz, 1H), 4.49 (br s, 2H), 3.84 (s, 3H); HRMS (m/z, relative intensity) 230 (M⁺, 100).


C. Ethyl 2-(4-amino-3-bromophenyl)acetate

Ethyl 2-(4-aminophenyl)acetate was used. Chromatography using elution with
5 ethyl acetate in hexanes [1:4] afforded the title compound (25%) as a light brown oil:
¹H NMR (CDCl₃) δ 7.33 (d, $J=2$ Hz, 1H), 7.02 (dd, $J=8$ and 2 Hz, 1H), 6.76 (dd, $J=8$ Hz, 1H), 4.11 (q, $J=7$ Hz, 2H), 3.45 (s, 2H), 1.23 (t, $J=7$ Hz, 3H); LRMS (m/z, relative intensity) 258 (M⁺, 100).

-28-

CLAIMS

1. A compound of the formula




10 where A represents a direct bond, C₁-C₄ alkyl, or C₁-C₄ alkenyl; n is 0, 1, or 2; R₁ is hydrogen, C₁-C₆ alkyl, aryl, C₁-C₃ alkylaryl, C₁-C₃ alkylheteroaryl, or -(CH₂)_mR₆; W, X, Y, and Z are each independently oxygen, sulfur, nitrogen or carbon, provided that at least one of W, X, Y or Z is nitrogen; R₂, R₃, R₄, and R₅ are each independently hydrogen, C₁-C₆ alkyl, aryl, C₁-C₃ alkylaryl, C₁-C₃ alkylheteroaryl, halogen, cyano, trifluoromethyl, nitro, -OR₇, -NR₇R₈, -(CH₂)_sOR₇, -SR₇, -SO₂NR₇R₈, -NR₇SO₂R₈,
 15 -NR₇CO₂R₈, -CONR₇R₈, or -CO₂R₇; one of R₂ and R₃, R₃ and R₄, or R₄ and R₅ may be taken together to form a five- to seven-membered alkyl ring, a six-membered aryl ring, a five- to seven-membered heteroalkyl ring having 1 heteroatom of N, O, or S, or a five- to six-membered heteroaryl ring having 1 or 2 heteroatoms of N, O, or S; R₆ is cyano, trifluoromethyl, or -OR₉; R₇, R₈, and R₉ are each independently hydrogen, C₁ to C₆ alkyl, -(CH₂)_mR₁₀, C₁ to C₃ alkylaryl, or aryl; R₇ and R₈ may be taken together to form a C₄-C₇ alkyl ring; R₁₀ is cyano, trifluoromethyl, or C₁-C₄ alkoxy; R₁₁ is hydrogen, -OR₁₂, or -NCHOR₁₂; R₁₂ is C₁ to C₆ alkyl, aryl, or C₁ to C₃ alkyl-aryl; m is 1, 2, or 3; s is 0, 1, 2, or 3; and the above aryl groups and the aryl moieties of the above alkylaryl groups are
 25 independently selected from phenyl and substituted phenyl, wherein said substituted phenyl may be substituted with one to three groups selected from C₁ to C₄ alkyl, halogen, hydroxy, cyano, carboxamido, nitro, and C₁ to C₄ alkoxy, and the pharmaceutically acceptable salts thereof.

2. A compound according to claim 1, wherein the compound of Formula

30 I is


-29-



3. A compound according to claim 2, wherein the compound is the cis epimer.

10 4. A compound according to claim 1 wherein A is a direct bond or -CH₂-; n is 1; R₁ is hydrogen, C₁-C₄ alkyl, or -CH₂CH₂OCH₃; Z is nitrogen; Y is carbon; W and X are each independently oxygen, sulfur, nitrogen or carbon; R₁₁ is hydrogen or -OCH₃.

5. A compound according to claim 4, wherein the compound of Formula I is



6. A compound according to claim 5, wherein the compound is the cis epimer.

7. A compound according to claim 1, said compound being selected from
 25 (R)-5-(4-benzyl-1,3-thiazol-2-yl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;
 (R)-5-(4-benzyl-1,3-thiazol-2-yl)-3-(pyrrolidin-2-ylmethyl)-1H-indole;
 (R)-5-(3-benzyl-1,2,4-oxadiazol-5-yl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-indole;
 (R)-5-(3-benzyl-1,2,4-oxadiazol-5-yl)-3-(pyrrolidin-2-ylmethyl)-1H-indole;
 (R)-5-(3-benzyl-1,2,4-oxadiazol-5-ylmethyl)-3-(N-methylpyrrolidin-2-ylmethyl)-1H-
 30 indole; and

(R)-5-(3-benzyl-1,2,4-oxadiazol-5-ylmethyl)-3-(pyrrolidin-2-ylmethyl)-1H-indole.

8. A pharmaceutical composition for treating a condition selected from hypertension, depression, anxiety, eating disorders, obesity, drug abuse, cluster

-30-


headache, migraine, pain, and chronic paroxysmal hemicrania and headache associated with vascular disorders comprising an amount of a compound according to claim 1 effective in treating such condition and a pharmaceutically acceptable carrier.

9. A pharmaceutical composition for treating disorders arising from deficient serotonergic neurotransmission comprising an amount of a compound according to claim 1 effective in treating such a disorder and a pharmaceutically acceptable carrier.

10. A method of treating a condition selected from hypertension, depression, anxiety, eating disorders, obesity, drug abuse, cluster headache, migraine, pain, and chronic paroxysmal hemicrania and headache associated with vascular disorders comprising administering to a mammal requiring such treatment an amount of a compound according to claim 1 effective in treating such condition.

11. A method for treating disorders arising from deficient serotonergic neurotransmission comprising administering to a mammal requiring such treatment an amount of a compound according to claim 1 effective in treating such condition.

12. A compound of the formula




where A represents a direct bond, C₁-C₄ alkyl, or C₁-C₄ alkenyl; n is 0, 1, or 2; W, X, Y, and Z are each independently oxygen, sulfur, nitrogen or carbon, provided that at least one of W, X, Y or Z is nitrogen; R₂, R₃, R₄, and R₅ are each independently hydrogen, C₁-C₆ alkyl, aryl, C₁-C₃ alkylaryl, C₁-C₃ alkylheteroaryl, halogen, cyano, trifluoromethyl, nitro, -O-, -NR₇R₈, -(CH₂)_mOR₇, -SR₇, -SO₂NR₇R₈, -NR₇SO₂R₈, -NR₇CO₂R₈, -CONR₇R₈, or -CO₂R₇; one of R₂ and R₃, R₃ and R₄, or R₄ and R₅ may be taken together to form a five- to seven-membered alkyl ring, a six-membered aryl ring, a five- to seven-membered heteroalkyl ring having 1 heteroatom of N, O, or S, or a five- to six-membered heteroaryl ring having 1 or 2 heteroatoms of N, O, or S; R₇ and R₈ are each independently hydrogen, C₁ to C₆ alkyl, -(CH₂)_mR₁₀, C₁ to C₃ alkylaryl, or aryl; R₇ and

-31-

R_8 may be taken together to form a C_4 - C_7 alkyl ring; R_{10} is cyano, trifluoromethyl, or C_1 - C_4 alkoxy; m is 1, 2, or 3; s is 0, 1, 2, or 3; R_{11} is hydrogen, $-OR_{12}$, or $-NHCOR_{12}$; R_{12} is C_1 to C_6 alkyl, aryl, or C_1 to C_3 alkyl-aryl; R_{13} is C_1 - C_6 alkyl, aryl, or alkylaryl; and the above aryl groups and the aryl moieties of the above alkylaryl groups are independently
 5 selected from phenyl and substituted phenyl wherein said substituted phenyl may be substituted with one to three groups selected from C_1 to C_4 alkyl, halogen, hydroxy, cyano, carboxamido, nitro, and C_1 to C_4 alkoxy.

13. A compound according to claim 12, wherein the compound of formula II is

10



15


II

14. A compound according to claim 13, wherein the compound is the cis epimer.

15. The compound of claim 12, wherein A is a direct bond or $-CH_2-$; n is 1, Z is nitrogen; Y is carbon; W and X are each independently oxygen, sulfur, nitrogen or carbon; R_{11} is hydrogen or $-OCH_3$.
 20

16. A compound according to claim 15, wherein the compound of formula II is

25




30

II

17. A compound according to claim 16, wherein the compound is the cis epimer.

-32-

18. A compound of the formula




where A represents a direct bond, C₁-C₄ alkyl, or C₁-C₄ alkenyl; n is 0, 1, or 2; W, X, Y, and Z are each independently oxygen, sulfur, nitrogen or carbon, provided that at least one of W, X, Y or Z is nitrogen; R₂, R₃, R₄, and R₅ are each independently hydrogen, C₁-C₆ alkyl, aryl, C₁-C₃ alkylaryl, C₁-C₃ alkylheteroaryl, halogen, cyano, trifluoromethyl, nitro, -OR₇, -NR₇R₈, -(CH₂)_sOR₇, -SR₇, -SO₂NR₇R₈, -NR₇SO₂R₈, -NR₇CO₂R₈, -CONR₇R₈, or -CO₂R₇; one of R₂ and R₃, R₃ and R₄, or R₄ and R₅ may be taken together to form a five- to seven-membered alkyl ring, a six-membered aryl ring, a five- to seven-membered heteroalkyl ring having 1 heteroatom of N, O, or S, or a five- to six-membered heteroaryl ring having 1 or 2 heteroatoms of N, O, or S; R₇ and R₈ are each independently hydrogen, C₁ to C₆ alkyl, -(CH₂)_mR₁₀, C₁ to C₃ alkylaryl, or aryl; R₇ and R₈ may be taken together to form a C₄-C₇ alkyl ring; R₁₀ is cyano, trifluoromethyl, or C₁-C₄ alkoxy; m is 1, 2, or 3; s is 0, 1, 2, or 3; R₁₁ is hydrogen, -OR₁₂, or -NHCOR₁₂; R₁₂ is C₁ to C₆ alkyl, aryl, or C₁ to C₃ alkyl-aryl; R₁₃ is C₁-C₆ alkyl, aryl, or alkylaryl; R₁₄ is halogen; and R₁₅ is -COCF₃, -SO₂CH₃, -SO₂Ph, or -CO₂C(CH₃)₃; and the above aryl groups and the aryl moieties of the above alkylaryl groups are independently selected from phenyl and substituted phenyl wherein said substituted phenyl may be substituted with one to three groups selected from C₁ to C₄ alkyl, halogen, hydroxy, cyano, carboxamido, nitro, and C₁ to C₄ alkoxy.

19. A compound according to claim 18, wherein the compound of formula III is

THIS PAGE BLANK (USPTO)


-33-



20. A compound according to claim 19, wherein the compound is the cis
 10 epimer.

21. The compound of claim 18, wherein A is a direct bond or -CH₂-; n is 1, Z is nitrogen; Y is carbon; W and X are each independently oxygen, sulfur, nitrogen or carbon; R₁₁ is hydrogen or -OCH₃.

22. A compound according to claim 21, wherein the compound of formula
 15 III is



23. A compound according to claim 22, wherein the compound is the cis
 25 epimer.

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC Int.Cl. 5 C07D417/14; A61K31/40; C07D413/14		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C07D	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	WO,A,9 118 897 (THE WELLCOME FOUNDATION LTD) 12 December 1991 see claims	1,8
P,A	EP,A,0 497 512 (MERCK SHARP & DOHME LTD.) 5 August 1992 see claims	1,8
A	EP,A,0 438 230 (MERCK SHARP & DOHME LTD.) 24 July 1991 cited in the application see claims	1,8
A	EP,A,0 313 397 (THE WELLCOME FOUNDATION LTD) 26 April 1989 cited in the application see claims	1,8
-/-		
¹⁰ Special categories of cited documents : ^{"A"} document defining the general state of the art which is not considered to be of particular relevance ^{"E"} earlier document but published on or after the international filing date ^{"L"} document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) ^{"O"} document referring to an oral disclosure, use, exhibition or other means ^{"P"} document published prior to the international filing date but later than the priority date claimed ^{"T"} later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention ^{"X"} document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step ^{"Y"} document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. ^{"A"} document member of the same patent family		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search 03 JUNE 1993		Date of Mailing of this International Search Report 21. 06. 93
International Searching Authority EUROPEAN PATENT OFFICE		Signature of Authorized Officer VAN BIJLEN H.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
P,A	WO,A,9 206 973 (PFIZER INC.) 30 April 1992 * complete document * ---	1,8
P,A	WO,A,9 213 856 (PFIZER INC.) 20 August 1992 see claims -----	1,8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 93/01667

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 10 and 11 are directed to a method of treatment of (diagnostic method practised on) the human/animal body the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims not searched : 10 and 11
Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9301667
SA 71115

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 03/06/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9118897	12-12-91	AU-A- 7957091 EP-A- 0486666	31-12-91 27-05-92
EP-A-0497512	05-08-92	AU-A- 1068092	06-08-92
EP-A-0438230	24-07-91	AU-A- 6944091 CN-A- 1053429	25-07-91 31-07-91
EP-A-0313397	26-04-89	AU-A- 2418188 JP-A- 1146882	27-04-89 08-06-89
WO-A-9206973	30-04-92	AU-A- 8950491 CN-A- 1062529	20-05-92 08-07-92
WO-A-9213856	20-08-92	AU-A- 1263792	07-09-92